



MAKHTESHIM
A G A N
of
NORTH AMERICA

BUMPER® 41.8EC
EPA Reg. No. 66222-42
FOR CONTROL OF ASIAN RUST ON SOYBEANS IN THE STATE OF MICHIGAN

**SECTION 18 EMERGENCY QUARANTINE EXEMPTION FOR THE USE OF BUMPER® 41.8EC
ON SOYBEANS FOR CONTROL OF ASIAN RUST (*Phakopsora pachyrhiza*) IN THE
STATE OF MICHIGAN**

EPA FILE SYMBOL: 04-MI-23
EXPIRES: 3-1-2007

ACTIVE INGREDIENT:	% BY WT.
Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]Methyl]-1 <i>H</i> -1,2,4-triazole.....	41.8%
INERT INGREDIENTS:	58.2%
	TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN
WARNING-AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- **FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**
- Follow all applicable directions, restrictions, Worker Protection Standards, and precautions on the EPA registered Bumper 41.8EC label.
- This label must be in the possession of the user at the time of pesticide application.

CROP, USE RATE, AND METHOD OF APPLICATION

If soybean rust is in the area or conditions are conducive to the disease, apply 4-8 fl. oz. per acre of Bumper 41.8EC. Repeat at 14-day intervals if conditions persist. It is important to protect the developing pod. Timing should be R1 (ca. 50 days after planting) to R5 depending on the conditions.

Coverage and penetration are important for best results. Bumper 41.8EC can be applied by either aerial or ground equipment. Use sufficient water volume to provide thorough and uniform plant coverage. When using aerial application, a minimum of 5 gallons per acre is recommended. With ground equipment, a minimum of 15 gallons per acre is recommended.

RESTRICTIONS

To avoid possible illegal residues:

1. Do not apply more than two applications of Bumper 41.8EC per acre per season, and no more than 12 fl. oz. of Bumper 41.8EC per acre per season.
2. Do not graze or feed soybean forage or hay.
3. Do not rotate to any crop intended for food, grazing, or any component of animal feed or bedding within 105 days of Bumper 41.8EC application unless the crop appears on the product label.
4. Do not apply later than soybean growth stage R5 (pod fill).
5. A maximum of two total applications using approved Section 18 products collectively are allowed under this soybean rust Section 18.
6. The REI is 24 hours.

Any adverse effects resulting for the use of Bumper 41.8EC under this emergency exemption must be immediately reported to the Michigan Department of Agriculture.

Makhteshim Agan of North America, Inc.
551 Fifth Ave., Suite 1100
New York, NY 10176

Bumper® is a registered trademark of Makhteshim Agan of North America, Inc.